

Is the MLB parameterization accurate enough to describe change in solar radiation with solar zenith angle?

Zhipeng Qu, Philippe Blanc, Mireille Lefèvre, Lucien Wald, Armel Oumbe

▶ To cite this version:

Zhipeng Qu, Philippe Blanc, Mireille Lefèvre, Lucien Wald, Armel Oumbe. Is the MLB parameterization accurate enough to describe change in solar radiation with solar zenith angle?. EGU General Assembly 2011, Apr 2011, Vienne, Austria. pp.EGU2011-9360-4. hal-00586438

HAL Id: hal-00586438 https://minesparis-psl.hal.science/hal-00586438v1

Submitted on 15 Apr 2011 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Is the MLB parameterization accurate enough to describe change in solar radiation with solar zenith angle?

Zhipeng Qu (1), Philippe Blanc (1), Mireille Lefèvre (1), Lucien Wald (1), and Armel Oumbe (2)
(1) MINES ParisTech, Centre for Energy and Processes, Sophia Antipolis, France (zhipeng.qu@mines-paristech.fr), (2)
German Remote Sensing Data Center, German Aerospace Center (DLR), Postfach 1116, D-82234 Wessling, Germany

A new direct method, Heliosat-4, is currently being developed by the MINES ParisTech and the German Aerospace Center (DLR), aiming at estimating surface downwelling solar irradiance (SSI). This method is composed of two parts: a clear-sky module and a cloud-ground module (Oumbe et al. 2009). The clear-sky model is based on the radiative transfer model (RTM) libRadtran fed with advanced products on atmosphere optical parameters derived from recent Earth Observation missions. However, running a RTM in an operational mode is computer resources and time consuming. Therefore, it will be very beneficial to use parameterization to reduce the number of runs of RTM.

Several parameterizations have been proposed in recent years, for example, the parameterization with water vapour density (Muller et al. 2009), with altitude (Gueymard, Thevenard 2009; Oumbe et al. 2009) and with sun zenith angle by using Modified Lambert-Beer function (MLB) (Muller et al. 2004, 2009). Qu et al. (2010) studied the MLB parameterization for beam and global irradiance. They proposed a 4-interval piecewise MLB which demonstrates better performances than the initial version based on a single interval of solar zenith angle. This piecewise MLB outperforms the current linear interpolation techniques. However, it is interesting to know whether the accuracy of this 4-piecewise MLB applied to global irradiance complies with standard performances expected for high quality pyranometers or for RTM. In order to assess it, we focused on two quality criteria: the systematic error (bias) and the percentile P95 of the absolute error. According to WMO standard for the measurement of radiation (WMO, 2008), the P95 should be less than 10 W/m2 (high quality). Myers et al. (2002) and Geuder et al. (2003) have compared different pyranometers. They wrote that as a whole a bias of 15 W/m2 is acceptable. Gueymard (2007) have compared results between different RTMs and estimated that a bias of 3 W/m2 is a proof of very high consistency between them. Therefore, we evaluated the performance of the 4-piecewise MLB by using two criteria: bias < 3 W/m2 and P95 < 10 W/m2.

We observed that the 4-piecewise MLB does not satisfy the criteria of P95 when the ground albedo is greater than 0.1. Therefore, we propose a 5-piecewise MLB in adding another fitting angle at 40. We observe that this 5-piecewise MLB demonstrates satisfactory performance and could be used accurately in the Heliosat-4 method to reduce the number of runs of the RTM.

Geuder, N., Trieb, F., Schillings, C., Meyer, R., Quaschning, V., Comparison of different methods for measuring solar irradiation data, 3rd International Conference on Experiences with Automatic Weather Stations, 19th-21st of February 2003, Torremolinos, Spain.

Gueymard, C. A., Prediction and validation of cloudless shortwave solar spectra incident on horizontal, tilted, or tracking surfaces, Solar energy, vol. 82, n°3, pp. 260-271, 2007.

Gueymard, C.A. and Thevenard, D.: Monthly average clear-sky broadband irradiance database for worldwide solar heat gain and building cooling load calculations. Solar Energy, vol. 83, 1998-2018, 2009.

Mueller, R., Dagestad, K.F., Ineichen, P., Schroedter, M., Cros, S., Dumortier, D., Kuhlemann, R., Olseth, J.A., Piernavieja, G., Reise, C., Wald, L., and Heinnemann, D.: Rethinking satellite based solar irradiance modelling - The SOLIS clear sky module, Remote Sensing of Environment, 91(2), 160-174, 2004.

Mueller, R.W., Matsoukas, C., Gratzki, A., Behr, H.D., and Hollmann, R.: The CM-SAF operational scheme for the satellite based retrieval of solar surface irradiance – A LUT based eigenvector hybrid approach, Remote Sensing of Environment, 113, 1012-1024, 2009.

Myers, D. R., Stoffel, T. L., Reda, I., Wilcox, S. M., Andreas, A. M., Recent progress in reducing the uncertainty in and improving pyranometer calibrations, Journal of Solar Energy Engineering, V.124, 44-50,2002.

Oumbe, A., Blanc, Ph., Ranchin, T., Schroedter-Homscheidt, M., and Wald, L.: A new method for estimating solar

energy resource. In Proceedings of the ISRSE 33, held in Stresa, Italy, 4-9 May 2009. Published by Joint Research Center, Ispra, Italy, USBKey, paper 773, 2009.

Qu, Z., Blanc, Ph., Lefèvre, M., Wald, L., Oumbe, A., Study of the MLB parameterisation for change in surface solar irradiance with sun zenith angle in clear sky, (submitted to Advances in Science and Research, Dec. 2010) World Meteorological Organization (WMO), Guide to Meteorological Instruments and Methods of Observation, Seventh edition, 2008.