



HAL
open science

Which effective solar zenith angles to use for hourly irradiation measurements?

Philippe Blanc, Lucien Wald

► To cite this version:

Philippe Blanc, Lucien Wald. Which effective solar zenith angles to use for hourly irradiation measurements?. 15th EMS / 12th ECAM, EMS European Meteorological Society, Sep 2015, Sofia, Bulgaria. pp.318. hal-01197035

HAL Id: hal-01197035

<https://minesparis-psl.hal.science/hal-01197035>

Submitted on 21 Sep 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Which effective solar zenith angles to use for hourly irradiation measurements?

Philippe Blanc and Lucien Wald

MINES ParisTech - PSL Research University, Sophia Antipolis cedex, France (lucien.wald@mines-paristech.fr)

Time-series of in situ or satellite-derived measurements of hourly irradiation on horizontal surface are increasingly available. In many cases, such data are input to numerical procedures that can be performed only if a solar zenith angle can be associated to each measurement. Currently, only time stamp is given for each measurement; no angle is given. The solar zenith angle is greatly varying within 1 h and cannot be considered as constant. In this context, there is a practical request from companies, academics, researchers...: what is the best practice for computing this angle? The work presented here compares the performances of several existing practices and makes recommendations keeping in mind the practical aspects faced by practitioners, companies, academics, researchers... High-quality 1 min measurements of global, diffuse and direct collected by the BSRN station in Carpentras in France, are aggregated to yield hourly irradiation, and used to assess the performances. It is found that several practices may be used to compute in a satisfactory way the effective solar zenith angle. The most accurate results are obtained if one may obtain corresponding irradiation values that should be observed under clear-sky conditions. The McClear Service available at <http://www.soda-pro.com> is one Web service providing such estimates of cloud-free irradiation for several summarizations, from 1 min to 1 month. If one may not obtain such cloud-free estimates, then the most accurate results are obtained from using irradiation at the top of atmosphere. Performances show a tendency to decrease during sunrise and sunset hours.