

Flint in its diverse natural occurrences: geo--tools for a better definition of the sourcing of secondary outcrops

Paul Fernandes, Michel Piboule, Médard Thiry, Céline Bressy-Leandri, François-Xavier Le Bourdonnec, Pascal Tallet, Jean--paul Raynal

▶ To cite this version:

Paul Fernandes, Michel Piboule, Médard Thiry, Céline Bressy-Leandri, François-Xavier Le Bourdonnec, et al.. Flint in its diverse natural occurrences: geo--tools for a better definition of the sourcing of secondary outcrops. On the Rocks', Xth International Symposium on Knappable Materials., Sep 2015, Barcelona, Spain. hal-01236679

HAL Id: hal-01236679 https://minesparis-psl.hal.science/hal-01236679

Submitted on 2 Dec 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Flint in its diverse natural occurrences: geo-tools for a better definition of the sourcing of secondary outcrops.

Paul Fernandes ¹⁻², Michel Piboule ³, Médard Thiry ⁴, Céline Leandri ⁵⁻⁶, François-Xavier Le Bourdonnec ⁷, Pascal Tallet ¹, Jean-Paul Raynal ^{2,8}

Precise identification of siliceous geo-resources used during prehistory still poses many problems, and archaeologists make ever-increasing demands for this data. The purpose is to provide a database containing an exact and descriptive identity for each different type of geological flint found within a region. The database is being developed for the administrative regions of Rhône-Alpes, Auvergne, Aquitaine and Provence-Alpes-Côte d'Azur. It will contribute to an accurate understanding of precise movements exchange of raw materials and therefore, to the behaviour of prehistoric groups.

Until now, the problem has been viewed in reverse; by trying to revert to the identity of the original material through study of the found object. Such a procedure follows an irrelevant path because it is first necessary to characterize raw materials in their primary position and secondly to make an appraisal of the epigenesis of the flint recovered from secondary sources.

The parameters we have chosen to characterize are the mineralogical composition (by optical microscopy, SEM, microprobe, cathodo-luminescence), microfacies characteristics (identified during microscopy and SEM image analysis), porosity measurements (by image analysis and porosimeter), and the presence and distribution of major and trace elements (using ICP, LA-ICP-MS, XRF, PIXE, Raman and SEM-EDS) at the surface or in the cracks in the matrix. Following the collection of the information, a statistical evaluation necessary to produce a reliable flint ID. Consideration of all varieties of the same material in its different states, coming from all known primary and secondary sources sets the study of archaeological samples on a solid foundation.

¹SARL Paléotime, 6173 rue Jean Séraphin Achard Picard, F-38350 Villard-de-Lans, France,

² Université de Bordeaux, CNRS UMR 5199 PACEA, PPP, bâtiment B18, allée Geoffroy Saint-Hilaire, CS 50023, 33615 Pessac cedex, France

³ 11. 21 rue General Ferrié. 38100 Grenoble

⁴ Mines-Paris-Tech, Géosciences, 35 rue ST Honoré, 77305 Fontainebleau, France.

⁵ Aix Marseille Université, CNRS, MCC, LAMPEA UMR 7269

⁶ Ministère de la Culture et de la Communication, DRAC de Corse, 20704 Ajaccio, France.

⁷ Université de Bordeaux Montaigne, Department of Art History and Archaeoloigy, Pessac, France

⁷8Departement of Human Evolution, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6, 04103 Leipzig, Allemagne.