

Convergence rate in the rough donsker theorem Laure Coutin, Laurent Decreusefond

▶ To cite this version:

Laure Coutin, Laurent Decreuse
fond. Convergence rate in the rough donsker theorem. 2017. hal
- $01551694\mathrm{v1}$

HAL Id: hal-01551694 https://minesparis-psl.hal.science/hal-01551694v1

Preprint submitted on 30 Jun 2017 (v1), last revised 13 Jun 2018 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CONVERGENCE RATE IN THE ROUGH DONSKER THEOREM

L. COUTIN AND L. DECREUSEFOND

ABSTRACT. We make precise the rate of convergence in the Lamperti's improvement of the Donsker Theorem on the topology of Hölder continuous functions. Then, we establish that the same rate of convergence holds true for the convergence of the enhanced, in the sense of rough paths theory, random walk towards the enhanced Brownian motion.

1. INTRODUCTION

The Donsker theorem says that a random walk

$$X^{m}(t) = \frac{1}{\sqrt{m}} \sum_{k=1}^{\lfloor mt \rfloor} X_{k}$$

where the X_k 's are independent, identically distributed random variables with mean 0 and variance 1, converges in a functional space to the Brownian motion B. In the original version (see [8]), the convergence was proved to hold in the space of continuous functions. The first evolution was the paper of Lamperti [13], which proved the convergence in Hölder spaces. Namely, he stated that if the increments of the X_k 's are *p*-integrable then the random walk X^m converges to B in $\operatorname{Hol}(1/2 - 1/p)$. The higher the integrability, the stronger the topology. There are numerous other extensions which can be made to the Donsker theorem. In the 90s, Barbour [2] estimated the rate of convergence in the space C of continuous functions on [0, 1] equipped with a stronger topology than the usual sup-norm topology. He proved that, provided that $\mathbf{E} [|X_k|^3]$ is finite, then

$$\sup_{|F|_M \le 1} \mathbf{E} \left[F(X^m) \right] - \mathbf{E} \left[F(B) \right] \le c \frac{\log m}{\sqrt{m}},$$

where M is roughly speaking, the set of thrice Fréchet differentiable functions on \mathcal{C} with bounded derivatives and $|F|_M$ is a function of the supremum of D^iF , $i = 0, \dots, 3$ over \mathcal{C} . The strategy is to compare X^m with B^m , the affine interpolation, of mesh 1/m, of the Brownian motion and then to compare B^m with B. The latter comparison is a sample-path comparison since the two processes live in the same probability space. This is the part which yields the log m factor. The former comparison is done via the Stein's method in finite dimension. The rate of this convergence is as usual (see [3, 14]) for Gaussian limits, of the order of $m^{-1/2}$.

²⁰⁰⁰ Mathematics Subject Classification. 60F17.

Key words and phrases. Donsker theorem, rough paths, Stein method.

In [5], we quantified the rate of convergence of X^m towards B in Besov-Liouville spaces ([15]), which are one scale of fractional Sobolev spaces. The spaces we considered were not included in Hölder spaces but the method could easily be adapted to obtain convergence rate in Hölder spaces. A slight adaptation of the proofs developed in this paper would show that under the same hypothesis as in [2],

$$\sup_{|F|_L \le 1} \mathbf{E} \left[F(X^m) \right] - \mathbf{E} \left[F(B) \right] \le cm^{2\beta - 1}$$

for any $\beta \in (0, 1/2)$ and L is the set of thrice Fréchet differentiable functions on $W_{\beta,3}$ (see below) with bounded derivatives equipped with the norm which is the sum of the sup-norm of F and of its first three derivatives.

Note that in both [2] and [5], an higher integrability of the X_k 's would not improve the convergence rates but would give more flexibility on the choice of the topology in which the convergence holds: The higher the integrability, the higher the Hölder exponent may be chosen.

In [11, Theorem 13.3.3], Friz and Victoir essentially showed that a Lamperti's like result holds for the convergence of the enriched random walk in the sense of rough path (see the definition below) to the enriched Brownian motion.

The motivation of this paper was to quantify the rate of this convergence in rough-paths sense. It raises several problems. The limiting process is no longer a Gaussian process: The Lévy area of a Brownian motion is not Gaussian. Hence we cannot expect to have a direct application of the Stein's method. However, there is no more randomness in the Lévy area that there is in the Brownian motion itself: The Lévy area is adapted to the filtration generated by the underlying Brownian motion. Saying that has two consequences. First, that the probability space we have to consider depends only on the Brownian motion. Second, that we have to find functional spaces for which the map which sends a Brownian motion to its Lévy area is not only continuous but also Lipschitz or at least locally Lipschitz. As mentioned in [12], the Besov-Liouville spaces are not well fitted to deal with the iterated integral processes we encounter in rough-paths theory. It is much better to work with the Slobodetsky scale of fractional Sobolev spaces.

In order to avoid some complicated calculations in infinite dimensional spaces, the idea is then to go back to the approach of [2], comparing the random walk with the affine interpolation of the Brownian motion in the Slobodetsky scale of fractional Sobolev spaces. This can be done by an application of the Stein's method in finite dimension. Then, we extend the norm on these spaces to take into account the iterated integrals of the enriched paths and show that the so-called signature map (see below) is Lipschitz continuous. The final result is obtained by considering the known distance between the enriched affine interpolation and the enriched Brownian motion in fractional Sobolev spaces.

Besides the definition of the convenient functional spaces, one of our contribution, interesting in its own, is an improvement of the Stein method. Actually, Stein's method for Gaussian limits in dimension 1 can be applied to C_b^1 functions but when the limit is in dimension 2 or more, the standard procedure works only for functions thrice differentiable with bounded derivatives. We show here that we can work with Lipschitz functions in any dimension by splitting the integral of the semi-group into two parts.

Rough paths theory is essentially a deterministic theory, it is therefore tempting to make the estimate we need in a deterministic setting and then to take the expectation of these bounds. This turns out to be a misleading approach. For instance, consider a sequence of centered, independent identically distributed random variables $(X_n, n \ge 1)$ and let $S_n = \sum_{i=1}^n X_i$. If we evaluate the *p*-th moment of S_n with Hölder inequality, we get that this moment is bounded by a constant times m^p . But if we use, as in the sequel, the Burkholder-Davis-Gundy inequality for discrete time martingale, we get an upper-bound proportional to $m^{p/2}$. This martingale argument which is implicitly used in [13] is the key to our work. For the sake of simplicity, this implies to separate the treatment of the symmetric and anti-symmetric parts of the signature. However, this is of no real importance since, as detailed below, the symmetric part of the signature can be handled as a classical \mathbf{R}^d -valued process.

This paper is organized as follows: In Section 2, we give the necessary notions about fractional Sobolev spaces and rough-paths theory. We also give a detailed proof of Lamperti's result in the fractional Sobolev spaces scale for further use and comparison. In Section 3, we then define the Kolmogorov-Rubinstein distance and show that this distance between the random walk and the affine interpolation of the Brownian motion can be reduced to a problem in finite dimension, should we consider a special set of Lipschitz functions. In Section 4, we then present our development of the Stein-Dirichlet method to estimate this distance.

2. Preliminaries

2.1. Fractional Sobolev spaces. As in [6, 12], we consider the fractional Sobolev spaces $W_{\eta,p}$ defined for $\eta \in (0, 1)$ and $p \geq 1$ as the the closure of \mathcal{C}^1 functions with respect to the norm

$$|f|_{\eta,p}^p = \int_0^1 |f(t)|^p \, \mathrm{d}t + \iint_{[0,1]^2} \frac{|f(t) - f(s)|^p}{|t - s|^{1+p\eta}} \, \mathrm{d}t \, \mathrm{d}s.$$

For $\eta = 1$, $W_{1,p}$ is the completion of \mathcal{C}^1 for the norm:

$$|f|_{\eta,p}^p = \int_0^1 |f(t)|^p \, \mathrm{d}t + \int_0^1 |f'(t)|^p \, \mathrm{d}t.$$

They are known to be Banach spaces and to satisfy the Sobolev embeddings [1, 10]:

$$W_{\eta,p} \subset \operatorname{Hol}(\eta - 1/p)$$
 for $\eta - 1/p > 0$

and

$$W_{\eta,p} \subset W_{\gamma,q}$$
 for $1 \ge \eta \ge \gamma$ and $\eta - 1/p \ge \gamma - 1/q$.

As a consequence, since $W_{1,p}$ is separable (see [4]), so does $W_{\eta,p}$.

2.2. Rough paths. We give a quick introduction to the rough-paths theory. For details, we refer to the monograph [11]. Consider $T^2(\mathbf{R}^d)$, the graded algebra of step two:

$$T^2(\mathbf{R}^d) = \mathbf{R} \oplus \mathbf{R}^d \oplus (\mathbf{R}^d \otimes \mathbf{R}^d).$$

We endow $T^2(\mathbf{R}^d)$ with an algebra structure $(+, ., \otimes)$ where for all (w_0, w_1, w_2) , $(z_0, z_1, z_2) \in T^2(\mathbf{R}^d), \ \lambda \in \mathbf{R}$

$$(w_0, w_1, w_2) + (z_0, z_1, z_2) = (w_0 + z_0, w_1 + z_1, w_2 + z_2)$$
$$\lambda.(w_0, w_1, w_2) = (\lambda w_0, \lambda w_1, \lambda w_2)$$

 $(w_0, w_1, w_2) \otimes (z_0, z_1, z_2) = (w_0 z_0, w_0 z_1 + z_0 w_1, w_0 z_2 + z_0 w_2 + w_1 \otimes z_1).$

Introduce the projection maps: For i = 0, 1, 2

$$\pi_i : T^2(\mathbf{R}^d) \longrightarrow (\mathbf{R}^d)^{\otimes i}$$
$$(w_0, w_1, w_2) \longmapsto w_i,$$

The set

$$1 + t^{2}(\mathbf{R}^{d}) = \{ w \in T^{2}(\mathbf{R}^{d}), \, \pi_{0}(w) = 1 \}$$
$$= \left\{ g = (1, w_{1}, w_{2}), \, (w_{1}, w_{2}) \in \mathbf{R}^{d} \oplus (\mathbf{R}^{d} \otimes \mathbf{R}^{d}) \right\}$$

is a Lie group with respect to the tensor multiplication \otimes , [11, Prop. 7.17]. Note that

$$(1, w_1, w_2)^{-1} = (1, -w_1, -w_2 + w_1 \otimes w_1)$$

As usual, a Lie group, like $1 + t^2(\mathbf{R}^d)$, leads to a Lie algebra when equipped with notions of product and commutator. Here, the Lie algebra is $(t^2(\mathbf{R}^d), +, .)$ with product \otimes and commutator

$$[g,w] = g \otimes w - w \otimes g = 0 \oplus (\pi_1(g) \otimes \pi_1(w) - \pi_1(w) \otimes \pi_1(g)),$$

for any $w, g \in t^2(\mathbf{R}^d).$

The exponential and logarithm maps are useful to go back and forth between $t^2(\mathbf{R}^d)$ and $1 + t^2(\mathbf{R}^d)$:

$$\exp : t^{2}(\mathbf{R}^{d}) \longrightarrow 1 + t^{2}(\mathbf{R}^{d})$$
$$w \longmapsto 1 + w + \frac{1}{2}(\pi_{1}w)^{\otimes 2}$$

and

$$\log : 1 + t^2(\mathbf{R}^d) \longrightarrow t^2(\mathbf{R}^d)$$
$$(1+w) \longmapsto w - \frac{1}{2}(\pi_1 w)^{\otimes 2}.$$

We denote by Σ the set of finite partitions $\sigma = \{t_1, \dots, t_n\}$ of [0, 1]. A continuous path z from [0, 1] into \mathbf{R}^d is said to have 1-finite variation whenever

$$\sup_{\sigma = \{t_1, \cdots, t_n\} \in \Sigma} \sum_{i=1}^{n-1} |z_{t_{i+1}} - z_{t_i}| < \infty.$$

The set of such functions equipped with this quantity as a norm is denoted by $C^{1-\text{var}}$.

Definition 2.1. The step-2 signature of $z \in C^{1-\text{var}}$ is given by:

$$S_2(z) : [0,1] \longrightarrow 1 + t^2(\mathbf{R}^d)$$
$$t \longmapsto \left(1, \ z_t - z_0, \ \int_0^t (z_s - z_0) \otimes dz_s\right).$$

The free nilpotent group of order 2, $G^2(\mathbf{R}^d)$, is the closed subgroup of $1 + t^2(\mathbf{R}^d)$ defined by

$$G^{2}(\mathbf{R}^{d}) = \{S_{2}(z), z \in C^{1-Var}\}.$$

We also consider $\check{G}_2(\mathbf{R}^d)$ (\check{G}_2 for short since d is fixed), the image of $G_2(\mathbf{R}^d)$ by the logarithm map. For $z \in C^{1-\text{var}}$, this corresponds to consider only the anti-symmetric part of $\pi_2(S_2(z))$:

$$\tilde{\mathbf{S}}_{2}(z) : [0,1] \longrightarrow t^{2}(\mathbf{R}^{d})
t \longmapsto \left(z_{t} - z_{0}, \int_{0}^{t} \left[(z_{s} - z_{0}), \mathrm{d}z_{s} \right] \right).$$

Remark 1. Denote by $(e_i, 1 \leq i \leq d)$ the canonical basis of \mathbf{R}^d , so that $(e_i \otimes e_j, 1 \leq i, j \leq d)$ is the canonical basis of $\mathbf{R}^d \otimes \mathbf{R}^d$. If

$$z(t) = \sum_{i=1}^{d} \sum_{k=1}^{m} z_{ik} h_k(t) e_i$$

where (h_1, \dots, h_m) are elements of $C^{1-\text{var}}$, we have

(1)
$$\log S_2(z)(t) = \left(z_t - z_0, \sum_{1 \le i < j \le d} \sum_{1 \le k < l \le m} z_{ik} z_{jl} \left(\int_0^t h_k(s) \, \mathrm{d}h_l(s) - \int_0^t h_l(s) \, \mathrm{d}h_k(s)\right) \, [e_i, e_j]\right).$$

For the sake of notations, we set $\mathcal{A} = \{1, \dots, d\} \times \{1, \dots, m\}$ and define the \prec relation by:

$$a = (a_1, a_2) \prec b = (b_1, b_2) \iff (a_1 < b_1) \text{ and } (a_2 < b_2).$$

With these notations, Eqn. (1) then becomes

$$\log S_2(z)(t) = \left(z_t - z_0, \\ \sum_{a \prec b} z_a z_b \left(\int_0^t h_{a_2}(s) \, \mathrm{d}h_{b_2}(s) - \int_0^t h_{b_2}(s) \, \mathrm{d}h_{a_2}(s) \right) \, [e_{a_1}, e_{b_1}] \right).$$

The group $G^2(\mathbf{R}^d)$ has the structure of a sub-Riemannian manifold. We will not dwell into the meanders of this very rich but intricate structure. It suffices to say that we can proceed equivalently by considering usual norms as follows.

For $\alpha \in (0,1)$, a path $w = 1 \oplus w_1 \oplus w_2$ is said to be α -Hölder whenever

$$\rho_{\alpha}(w) = \max\left(\sup_{s \neq t} \frac{|w_1(t) - w_1(s)|}{|t - s|^{\alpha}}, \sup_{s \neq t} \frac{|\pi_2(w(s)^{-1} \otimes w(t))|^{1/2}}{|t - s|^{\alpha}}\right) < \infty.$$

Note that for $z \in C^{1-\operatorname{var}}$,

$$\rho_{\alpha}(S_{2}(z)) = \max\left(\sup_{s \neq t} \frac{|z(t) - z(s)|}{|t - s|^{\alpha}}, \sup_{s \neq t} \frac{\left|\int_{s}^{t} (z_{r} - z_{s}) \otimes dz_{r}\right|^{1/2}}{|t - s|^{\alpha}}\right).$$

Definition 2.2. We denote by $H_{\alpha}(t^2(\mathbf{R}^d))$, the vector space of paths w from [0,1] into $t^2(\mathbf{R}^d)$ such that $\rho_{\alpha}(w)$ is finite. It is equipped with the homogeneous norm: For w and v in $H_{\alpha}(t^2(\mathbf{R}^d))$

$$|w - v|_{H_{\alpha}(t^{2}(\mathbf{R}^{d}))} = \rho_{\alpha}(w - v) = \rho_{\alpha}((w_{1} - v^{(1)}) \oplus (w_{2} - v^{(2)})).$$

Unfortunately, as mentioned in [11, Chapter 8.3], this metric space is complete but not separable, which is unacceptable for our purpose (see Definition 3.1 and the remark below). We thus introduce fractional Sobolev spaces as in [12].

Definition 2.3. For any $\eta \in (0, 1)$, any $p \ge 2$, $\tilde{G}_2 W_{\eta,p}$ is the vector space of paths w from [0, 1] into $t^2(\mathbf{R}^d)$ such that

$$|w_1|_{\eta,p}^p + \iint_{[0,1]^2} \frac{\left|\frac{\pi_2[w_s^{-1}, w_t]\right|^{p/2}}{|t-s|^{1+\eta p}} \, \mathrm{d}s \, \mathrm{d}t < \infty.$$

The distance on $\tilde{G}_2 W_{\eta,p}$ is defined by

$$|w - v|_{\tilde{G}_{2}W_{\eta,p}} = |\pi_{1}(w) - \pi_{1}(v)|_{W_{\eta,p}} + \left(\iint_{[0,1]^{2}} \frac{\left|\pi_{2}[w_{s}^{-1}, w_{t}] - \pi_{2}[v_{s}^{-1}, v_{t}]\right|^{p/2}}{|t - s|^{1 + \eta p}} \, \mathrm{d}s \, \mathrm{d}t \right)^{1/p}$$

Following [12], we know that $\check{G}_2 W_{\eta,p}$ is a Banach space included into $H_{\alpha}(t^2(\mathbf{R}^d))$ for $\alpha = \eta - 1/p$, provided $\alpha > 0$.

Lemma 2.1. For any $\eta \in (0,1)$, any $p \ge 2$, $\check{G}_2 W_{\eta,p}$ is separable.

Proof. Consider the map κ defined as

$$\kappa : ([0,1] \to t_2(\mathbf{R}^d)) \longrightarrow ([0,1]^2 \to t_2(\mathbf{R}^d))$$
$$w \longmapsto \left((s,t) \mapsto \left(\pi_1(w_t) - \pi_1(w_s), \, \pi_2[w_s^{-1}, \, w_t] \right) \right).$$

Consider the measure $d\mu_{\eta,p}(s,t) = |t-s|^{-1-\eta p} ds dt$. Then,

(2)
$$|w|_{\check{G}_2W_{\eta,p}} = |\pi_1 \circ \kappa(w)|_{L^p(\mu_{\eta,p})} + |\pi_2 \circ \kappa(w)|_{L^{p/2}(\mu_{2\eta,p/2})}^{1/2}.$$

For any $p \geq 1$ and $\eta \in (0,1)$, $L^p(\mu_{\eta,p})$ is isometrically isomorphic to $L^p(\mathrm{d}s \, \mathrm{d}t)$ hence it is separable. This entails that $E = L^p(\mu_{\eta,p}) \times L^{p/2}(\mu_{2\eta,p/2})$ is separable. Equation (2) means that the application T which maps $w \in \check{G}_2 W_{\eta,p}$ to the couple $(\pi_1 \circ \kappa(w), \pi_2 \circ \kappa(w)) \in E$, is an isometry. Thus $\check{G}_2 W_{\eta,p}$ is isometrically isomorphic to a closed subspace of the separable space E, hence it is separable. \Box

2.3. **Donsker-Lamperti theorem.** For the sake for completeness and for further comparison, we give the proof of the Donsker-Lamperti theorem in the scale of fractional Sobolev spaces, which induces the convergence in Hölder spaces.

Definition 2.4. The random walk associated to the sequence $(X_k, k \ge 1)$ is defined by

$$X^{m}(t) = \sqrt{m} \sum_{k=1}^{m} X_{k} r_{k}^{m}(t) = \sum_{k=1}^{m} X_{k} h_{k}^{m}(t)$$

where

$$r_k^m(t) = \int_0^t \mathbf{1}_{((k-1)/m, k/m]}(s) \, \mathrm{d}s \text{ and } h_k^m = \sqrt{m} \, r_k^m.$$

Theorem 2.2. If for any $k \ge 1$, X_k belongs to L^p for some $p \ge 2$, then there exists c > 0 such that

(3)
$$\sup_{m \ge 1} \frac{\mathbf{E}\left[\left|\sum_{k=1}^{m} X_k \sqrt{m} \left(r_k^m(t) - r_k^m(s)\right)\right|^p\right]}{|t - s|^{p/2}} < c \mathbf{E}\left[|X_1|^p\right]$$

Proof. For $0 \le s < t \le 1$ fixed, the discrete time process

$$Y_m^{st} = \sum_{k=1}^m X_k (r_k^m(t) - r_k^m(s))$$

is a martingale with respect to the filtration $\mathcal{F}_n = \sigma(X_k, 1 \leq k \leq m)$. The Burkholder-Davis-Gundy [17] entails that

$$m^{p}\mathbf{E}\left[\left|\sum_{k=1}^{m} X_{k}\left(r_{k}^{m}(t)-r_{k}^{m}(s)\right)\right|^{p}\right] \leq c \, m^{p/2}\mathbf{E}\left[\left|\sum_{k=1}^{m} X_{k}^{2}\left(r_{k}^{m}(t)-r_{k}^{m}(s)\right)^{2}\right|^{p/2}\right].$$

If $|t-s| \leq 1/m$, there is at most two values of k such that $r_k^m(t) - r_k^m(s)$ is not zero. Furthermore,

$$|r_k^m(t) - r_k^m(s)| \le |t - s|$$

hence

$$|r_k^m(t) - r_k^m(s)|^2 \le m^{-1}|t - s|.$$

In this situation,

$$\mathbf{E}\left[\left|\sum_{k=1}^{m} X_{k}^{2} \left(r_{k}^{m}(t) - r_{k}^{m}(s)\right)^{2}\right|^{p/2}\right] \leq c \, m^{-p/2} \, \mathbf{E}\left[|X_{1}|^{p}\right] \, |t - s|^{p/2},$$

so that (3) holds true for $|t-s| \leq 1/m$. For |t-s| > 1/m, we remark that $r_k^m(t) - r_k^m(s)$ is not null for at most [m(t-s)] + 2 values of k and since r_k^m is Lipschitz continuous, $|r_k^m(t) - r_k^m(s)| \leq 1/m$ for such value of k. Hence, by convexity inequality,

$$\mathbf{E}\left[\left|\sum_{k=1}^{m} X_{k}^{2} \left(r_{k}^{m}(t) - r_{k}^{m}(s)\right)^{2}\right|^{p/2}\right] \leq c \left(m|t-s|+2\right)^{p/2} m^{-p} \mathbf{E}\left[|X_{1}|^{p}\right]$$
$$\leq cm^{-p/2} \mathbf{E}\left[|X_{1}|^{p}\right] |t-s|^{p/2}.$$

Hence, (3) is true for $|t - s| \ge 1/m$.

It is then straightforward that we have:

Corollary 2.3. Assume that for any $k \ge 1$, X_k belongs to L^p for some $p \ge 2$. Then, for any $\eta < 1/2$,

$$\sup_{m\geq 1} \mathbf{E}\left[|X^m|_{W_{\eta,p}}^p\right] < \infty.$$

Proof. Actually, $(s,t) \mapsto |t-s|^{p/2}$ is $\mu_{\eta,p}$ -integrable provided that $p(1/2 - \eta) > 0$, i.e. $\eta < 1/2$.

Corollary 2.4 (Lamperti). Assume that for any $k \ge 1$, X_k belongs to L^p for some $p \ge 2$. Then, for any $1/p < \eta < 1/2$, the sequence $(X^m, m \ge 1)$ converges in distribution in $\operatorname{Hol}(\eta - 1/p)$ to B.

Proof. It is well-known that the finite dimensional distributions of X^m converge to that of B. From the previous lemma, we know that for any $0 < \eta < \zeta < 1/2$, there exists K such that

$$\sup_{m\geq 1} \mathbf{P}(|X^m|_{W_{\zeta,p}}) \geq K) \leq \eta.$$

The embedding of $W_{\zeta,p}$ into $W_{\eta,p}$ is compact, hence the sequence $(X^m, m \ge 1)$ is tight in $W_{\eta,p}$ hence convergent. The result follows by the continuous embedding of $W_{\eta,p}$ into $\operatorname{Hol}(\eta - 1/p)$.

3. Rate of convergence

3.1. Kolmogorov-Rubinstein distance. The proof of Lamperti's Theorem is given for one dimensional processes but it can be straightforwardly adapted to \mathbf{R}^d -valued random walks and Brownian motion: X^m becomes the \mathbf{R}^d -valued process

$$X^{m}(t) = \sqrt{m} \sum_{a \in \mathcal{A}} X_{a} r_{a_{2}}(t) e_{a_{1}} = \sum_{a \in \mathcal{A}} X_{a} h_{a}(t)$$

where $h_a(t) = \sqrt{m} r_{a_2}(t) e_{a_1}$ and $(X_a, a \in \mathcal{A})$ is a family of independent identically distributed random variables of mean 0 and variance 1. Furthermore, B is the *d*-dimensional Brownian motion:

$$B(t) = \sum_{i=1}^{d} B_i(t) e_i.$$

The enriched Brownian motion \mathbb{B} , is the $G_2(\mathbf{R}^d)$ -value process defined by

$$\mathbb{B}(t) = 1 \oplus B(t) \oplus \sum_{i,j=1}^{d} \int_{0}^{t} B_{i}(s) \circ \mathrm{d}B_{j}(s) \ e_{i} \otimes e_{j}, \text{ for any } t \in [0,1],$$

where the stochastic integrals are to be understood in the Stratonovitch sense. Theorem 13.32 of [11] says that $S_2(X^m)$ converges to \mathbb{B} in some Hölder type spaces. Our primary goal is to give the rate of this convergence. For, we need to define a distance between probability measures over Hölder spaces. There are several possibilities of such a definition, the best suited for an estimate by the Stein method is the Kolmogorov-Rubinstein distance: **Definition 3.1** (Kolmogorov-Rubinstein distance). For μ and ν two probability measures on a metric space (W, d_W) , their Kolmogorov-Rubinstein distance is given by

$$\operatorname{dist}_{\operatorname{KR}}(\mu, \nu) = \sup_{F \in \operatorname{Lip}(d_W)} \int_W F \, \mathrm{d}\mu - \int_W F \, \mathrm{d}\nu,$$

where

$$\operatorname{Lip}(d_W) = \Big\{ F : W \to \mathbf{R}, \ \forall w, v \in W, \ |F(w) - F(v)| \le d_W(w, v) \Big\}.$$

Theorem 11.3.3 of [9] states that the topology induced by this distance on the set of probability measures on W is the same as the topology of the convergence in laws whenever the metric space W is separable. Unfortunately, as we already mentioned, Hölder spaces are not separable, thus to have a meaningful result, we turn to work on fractional Sobolev spaces. It is of no importance since Sobolev embeddings ensure that convergence in fractional Sobolev spaces induces convergence in Hölder spaces as for the Lamperti Theorem.

Our new goal is then to estimate the Kolmogorov-Rubinstein distance in $G_2W_{\eta,p}$ between \mathbb{B} and $S_2(X^m)$. Remark that

$$\pi_2 S_2(X^m) = \pi_2 \check{S}_2(X^m) + 2 \sum_{a \in \mathcal{A}} (X_a^2 - 1) \int_0^t h_a^m(s) \otimes dh_a^m(s) + \sum_{a \in \mathcal{A}} h_a^m(t) \otimes h_a^m(t) = U_1^m + U_2^m + U_2^m.$$

On the other hand,

$$\mathbb{B}(t) = \sum_{1 \le i < j \le d} \left(\int_0^t B_i(s) \circ dB_j(s) - \int_0^t B_j(s) \circ dB_i(s) \right) [e_i, e_j] + 2 \sum_{i=1}^d \int_0^t B_i(s) dB_i(s) e_i \otimes e_i + \sum_{i=1}^d t e_i = \log \mathbb{B}(t) + U_2 + U_3.$$

where in U_2 , the stochastic integral is taken in the Itô sense. Direct computations show that $U_3^m - U_3$ tends to 0 as 1/m. The convergence of U_2^m towards U_2 is of the same kind as the standard Donsker theorem: It follows the same convergence rate as that of the convergence of X^m to B; a rate which is expected and which will turn out to be much slower than 1/m. In summary, the Kolmogorov-Rubinstein distance between \mathbb{B} and $S_2(X^m)$ has the same asymptotic behavior as the distance between $\log \mathbb{B}$ and $\check{S}_2(X^m)$. Our final objective is then to estimate the Kolmogorov-Rubinstein distance between the distributions of $\log \mathbb{B}$ and $\check{S}_2(X^m)$ in $\check{G}_2 W_{\eta,p}$.

3.2. Reduction to finite dimension. Should we follow the same procedure as the one we used in [5], we would face the same complications to compute the trace term in some infinite dimensional space. To circumvent this difficulty, we decompose the distance into two parts: Consider B^m the affine interpolation of B, we use the results of [11] to estimate the distance between $\check{S}_2(B^m)$ and $\log \mathbb{B}$ and then resort to the Stein's method in finite dimension to estimate the distance between $\check{S}_2(X^m)$ and $\check{S}_2(B^m)$. We can always write

$$\sup_{F \in \operatorname{Lip}(d_{\check{G}_{2}W_{\eta,p}})} \mathbf{E} \left[F(\log \mathbb{B}) \right] - \mathbf{E} \left[F(\check{S}_{2}(X^{m})) \right]$$

$$\leq \sup_{F \in \operatorname{Lip}(d_{\check{G}_{2}W_{\eta,p}})} \mathbf{E} \left[F(\log \mathbb{B}) \right] - \mathbf{E} \left[F(\check{S}_{2}(B^{m})) \right]$$

$$+ \sup_{F \in \operatorname{Lip}(d_{\check{G}_{2}W_{\eta,p}})} \mathbf{E} \left[F(\check{S}_{2}(B^{m})) \right] - \mathbf{E} \left[F(\check{S}_{2}(X^{m})) \right].$$

On the one hand, since $\log \mathbb{B}$ and $\check{S}_2(B^m)$ live on the same probability space, for $F \in \operatorname{Lip}(d_{\check{G}_2W_{\eta,p}})$, according to [11, Proposition 13.20], (4)

$$\mathbf{E}\left[F(\log \mathbb{B})\right] - \mathbf{E}\left[F(\check{\mathbf{S}}_{2}(B^{m}))\right] \leq \mathbf{E}\left[\left|\log \mathbb{B} - \check{\mathbf{S}}_{2}(B^{m})\right|_{\check{\mathbf{G}}_{2}W_{\eta,p}}\right] \leq c \, m^{-(1/2-\eta)}.$$

It remains to estimate

 $F \in I$

$$\sup_{F \in \operatorname{Lip}(d_{\tilde{G}_2W_{n,n}})} \mathbf{E} \left[F(\check{S}_2(B^m)) \right] - \mathbf{E} \left[F(\check{S}_2(X^m)) \right]$$

Theorem 3.1. If $p \ge 3$ and X_a belongs to L^p , for any $\eta \in (1/p, 1/2)$,

 $\sup_{F \in \operatorname{Lip}(\check{\mathbf{G}}_2 W_{\eta,p})} \mathbf{E} \left[F(\check{\mathbf{S}}_2(B^m)) \right] - \mathbf{E} \left[F(\check{\mathbf{S}}_2(X^m)) \right] \le c \, \|X_a\|_{L^p} \, m^{-1/2+\eta}.$

Proof of Theorem. According to Theorem 4.5,

$$\mathbf{E}\left[\left|P_{\tau_0}F(\check{S}_2(X^m) - F(\check{S}_2(X^m))\right|\right] \le c \, \|X_a\|_{L^p} \, m^{-1/2+\eta} \, \sqrt{1 - e^{-\tau_0}}.$$

Combine this upper-bound with (8) to obtain

$$\sup_{\text{Lip}(\check{G}_{2}W_{\eta,p})} \mathbf{E} \left[F(\check{S}_{2}(B^{m})) \right] - \mathbf{E} \left[F(\check{S}_{2}(X^{m})) \right]$$
$$\leq c \, m^{-1/2+\eta} \, \|X_{a}\|_{L^{p}} \left(\sqrt{1 - e^{-\tau_{0}}} + m^{2\eta} \, \int_{\tau_{0}}^{\infty} \frac{e^{-5\tau/2}}{1 - e^{-\tau/2}} \, \mathrm{d}\tau \right).$$

The bracketed term is decreasing with respect to τ_0 and tends to 1 as τ_0 goes to infinity, hence the result.

We now detail the proofs of the main estimates.

4. Improved Stein Method

4.1. Finite dimensional Gelfand triplet. The title of this section may seem pompous at first reading but the construction of the Wiener space we are working on is of crucial importance and not so usual. In view of the previous considerations, our computations are to be held in a finite dimensional space and Gelfand triplet (or abstract Wiener space) are mainly thought to be relevant for infinite dimensional spaces. Actually, our settings should reflect the fact that we are in some sense, projecting an abstract Wiener infinite dimensional space onto a finite dimensional Gelfand triplet.

Recall that $\mathcal{A} = \{1, \dots, d\} \times \{1, \dots, m\}$. Let \mathcal{V}_m be the finite dimensional vector space

$$\mathcal{V}_m = \operatorname{span}\{h_a^m, \, a \in \mathcal{A}\}$$

We put on \mathcal{V}_m the norm of $W_{\eta,p}$. We denote by W, the space \mathcal{V}_m equipped with this norm. To simplify the forthcoming computations, it is useful to introduce

$$g_a^m = \alpha_m^{-1} h_a^m = \alpha_m^{-1} \left(h_{a_2} \otimes e_{a_1} \right),$$

where the coefficient α_m is chosen so that $|g_a^m|_{W_{\eta,p}} = 1$.

Furthermore, consider the space H which is \mathcal{V}_m equipped with the scalar product:

$$(g_a^m, g_b^m)_H = \delta_{a,b}$$

Recall that

$$B^m = \sum_{a \in \mathcal{A}} B_a h_a^m = \alpha_m \sum_{a \in \mathcal{A}} B_a g_a^m.$$

Let \mathbf{P}^m be the distribution of B^m , it is a centered Gaussian probability measure on \mathcal{V}_m with covariance matrix α_m^2 Id. The triplet (H, W, \mathbf{P}^m) is a Gelfand triplet: \mathbf{P}^m is a Gaussian measure on (the Banach space) W, which contains (as a dense subset) the (Hilbert) space H and

$$\int_{W} e^{i\langle\eta,\omega\rangle_{W^*,W}} \,\mathrm{d}\mathbf{P}^m(\omega) = \exp(-\frac{1}{2}|\iota^*(\eta)|_H^2)$$

where

$$W^* \xrightarrow{\iota^*} H^* \simeq H \xrightarrow{\iota} W.$$

In a finite dimensional setting, all these spaces and their dual are isomorphic and isomorphic to \mathbf{R}^{dm} , but we cannot identify more than one space with its dual, hence the necessity to define precisely the Gelfand triplet.

4.2. Malliavin-Dirichlet structure. In particular, the Cameron-Martin theorem reads as:

(5)
$$\mathbf{E}\left[F(B^m + \alpha_m \sum_{a \in \mathcal{A}} y_a g_a^m)\right]$$
$$= \mathbf{E}\left[F(B^m) \exp\left(\alpha_m \langle y, B^m \rangle_H - \frac{\alpha_m^2}{2} |y|_H^2\right)\right].$$

We denote by ∇F the *H* valued gradient of *F* provided that it exists: For any $v \in \mathcal{V}_m$, for any $g \in$,

$$\langle \nabla F(v), g \rangle_{H,H} = \frac{d}{d\varepsilon} F(v + \varepsilon g) \bigg|_{\varepsilon=0}.$$

The space of twice differentiable functions on \mathcal{V}_m is denoted by $\mathcal{C}^2(\mathcal{V}_m)$. The Fréchet Laplacian is defined by:

$$\Delta F(v) = \operatorname{trace} \nabla^{(2)} F(v) = \sum_{a \in \mathcal{A}} \left\langle \nabla^{(2)} F(v), \, g_a^m \otimes g_a^m \right\rangle_{H^{\otimes 2}, H^{\otimes 2}}.$$

Let L be the operator defined on $\mathcal{C}^2(\mathcal{V}_m)$ by:

$$LF(v) = -\left\langle \nabla F(v), v \right\rangle_{H,H} + \alpha_m^2 \,\Delta F(v).$$

The Cameron-Martin theorem (see Eqn. (5)) yields an integration by parts formula on W which can be interpreted by saying that the distribution of B^m on \mathcal{V}_m is characterized by $\mathbf{E}[LF] = 0$ for any $F \in \mathcal{C}^2(\mathcal{V}_m)$. It induces an Ornstein-Uhlenbeck semi-group $(P_t, t \ge 0)$ which can be described by the Mehler formula:

$$P_t F(v) = \int_{\mathcal{V}_m} F(e^{-t}v + \beta_t w) \, \mathrm{d}\mathbf{P}^m(w),$$

where $\beta_t = (1 - e^{-2t})^{1/2}$.

Introduce for any $a \prec b \in \mathcal{A}$ and any $0 \leq s < t \leq 1$,

$$\check{g}_{a,b}^{m}(s,t) = \left\{ \int_{s}^{t} \left(g_{a_{2}}^{m}(r) - g_{a_{2}}^{m}(s) \right) \, \mathrm{d}g_{b_{2}}^{m}(r) - \int_{s}^{t} \left(g_{b_{2}}^{m}(r) - g_{b_{2}}^{m}(s) \right) \, \mathrm{d}g_{a_{2}}^{m}(r) \right\} \, [e_{a_{1}}, e_{b_{1}}].$$

Lemma 4.1. For any $\omega = (\omega_k, 1 \le k \le m)$, we have

$$\left|\sum_{k=1}^{m} \omega_k h_k^m\right|_{var,[s,t]} \le \sqrt{m} \sum_{k=1}^{m} |\omega_k| \left\{ (t-s) \land \frac{1}{m} \right\} \cdot$$

Moreover,

$$|\sum_{k=1}^{m} \omega_k h_k^m|_{\eta,p} \le c \, m^{-(1/2-\eta)}.$$

Proof. For $k \neq \ell$, $dh_k^m dh_\ell = 0$ hence their contribution to the overall variation are separate:

$$\|\sum_{k=1}^{m} \omega_k h_k^m\|_{\operatorname{var},[s,t]} = \sum_{k=1}^{m} |\omega_k| \|h_k^m\|_{\operatorname{var},[s,t]}.$$

It is easy to see that

$$\|h_k^m\|_{\operatorname{var},[s,t]} = \sqrt{m} \ (t \wedge \frac{k}{m} - s \vee \frac{k-1}{m}) \le \sqrt{m} \ \left\{ (t-s) \wedge \frac{1}{m} \right\} \cdot$$
roof is thus complete.

The proof is thus complete.

Lemma 4.2. For any $\tau > 0$, for any $v \in \mathcal{V}_m$, for any $F \in \operatorname{Lip}(\check{\mathrm{G}}_2 W_{n,p})$, we have

(6)
$$\left\langle \nabla^{(2)} P_{\tau} F(v), g_a^m \otimes g_a^m \right\rangle$$

= $\frac{e^{-3\tau/2}}{\beta_{\tau/2}^2} \iint F\left(v(\tau, y, z)\right)(y, g_a^m)_H(z, g_a^m)_H d\mathbf{P}^m(y) d\mathbf{P}^m(z),$

where

$$v(\tau, y, z) = e^{-\tau/2} (e^{-\tau/2} v + \beta_{\tau/2} y) + \beta_{\tau/2} z.$$

Moreover, there exists a constant c such that

(7)
$$\left| \left\langle \nabla^{(3)} P_{\tau} F(v), (g_a^m)^{\otimes(3)} \right\rangle \right|$$

 $\leq c \frac{e^{-5\tau/2}}{\beta_{\tau/2}^2} \left(\iint \left| \sum_{b \prec a} \left(v_b \check{g}_{a,b}^m \right)_{st} \right|^{p/2} d\mu_{\eta,p}(t,s) \right)^{1/p}$

Proof. Following [16, Lemma 4.5], we have

$$\left\langle \nabla^{(2)} P_{\tau} F(v), \, g_a^m \otimes g_a^m \right\rangle$$

= $\frac{e^{-\tau}}{\beta_{\tau/2}} \int \left\langle \nabla P_{\tau/2} F(e^{-\tau/2}v + \beta_{\tau/2}y), \, g_a^m \right\rangle \, (y, g_a^m)_H \, \mathrm{d}\mathbf{P}^m(y),$

where we use integration by parts to have the following representation of the first order gradient:

$$\langle \nabla P_{\tau/2} F(w), g_a^m \rangle = \frac{e^{-\tau/2}}{\beta_{\tau/2}} \int F(e^{-\tau/2}w + \beta_{\tau/2}z) (z, g_a^m)_H \, \mathrm{d}\mathbf{P}^m(z).$$

Plugging in this equation in the former yields (6). Moreover,

$$(v + \epsilon g_a^m)(\tau, y, z) = \epsilon e^{-\tau} g_a^m + v(\tau, y, z)$$

and

$$\begin{split} \left| \check{\mathbf{S}}_{2} \Big(\epsilon e^{-\tau} g_{a}^{m} + v(\tau, y, z) \Big) - \check{\mathbf{S}}_{2} \Big(v(\tau, y, z) \Big) \right|_{W_{\eta, p}} \\ & \leq \epsilon \, e^{-\tau} \left(1 + \left(\iint \left| \sum_{b \prec a} \left(v_{b} \, \check{g}_{a, b}^{m} \right)_{st} \right|^{p/2} \, \mathrm{d}\mu_{\eta, p}(t, s) \right)^{1/p} \right). \end{split}$$
ence, (7) holds true.

Hence, (7) holds true.

Lemma 4.3. Let $(U_a, a \in A)$ be a family of independent identically distributed random variables which belong to L^p . Then,

$$\mathbf{E}\left[\left|U_a \sum_{b \prec a} U_b \check{g}^m_{a,b}\right|_{W_{\eta,p}}^p\right] \le c \, \mathbf{E} \left[|U_a|^p\right]^2.$$

Proof. By independence and as in the proof of Theorem 2.2,

$$\mathbf{E}\left[\left|U_a\sum_{b\prec a}U_b\check{g}^m_{a,b}(s,t)\right|^p\right] \le c\,\mathbf{E}\left[\left|U_a\right|^p\right]\mathbf{E}\left[\left(\sum_{b\prec a}|U_b|^2|\check{g}^m_{a,b}(s,t)|^2\right)^{p/2}\right].$$

Since a is fixed and $b \prec a$, if $|t-s| \leq 1/m$, $\check{g}^m_{a,b}$ is not zero only for $b_2 = a_2 - 1$ and then, it is bounded by $|t-s|/\sqrt{m}$. If $|t-s| \geq 1/m$, $\check{g}^m_{a,b}$ is not zero for at most [dm|t-s|] values of b and then, each term is bounded by 1/m. Hence

$$\mathbf{E}\left[\left(\sum_{b\prec a} |U_b|^2 |\check{g}_{a,b}^m(s,t)|^2\right)^{p/2}\right] \le c \, m^{-p/2} (m|t-s|)^{p/2} \mathbf{E}\left[|U_b|^p\right].$$

The result follows by integration with respect to $\mu_{\eta,p}$.

4.3. Main computations. Recall that

$$X^m = \alpha_m \sum_{a \in \mathcal{A}} X_a g_a^m \text{ and } B^m = \alpha_m \sum_{a \in \mathcal{A}} B_a g_a^m.$$

and that we expect to estimate the supremum, for $F \in \text{Lip}(\dot{G}_2 W_{\eta,p})$, of

$$\mathbf{E}\left[F(\check{\mathbf{S}}_2B^m)\right] - \mathbf{E}\left[F(\check{\mathbf{S}}_2X^m)\right].$$

For the sake of notations, we set $\check{F} = F \circ \check{S}_2$. The Stein-Dirichlet representation formula (see [7]) then stands that

$$\mathbf{E}\left[\check{F}(B^{m})\right] - \mathbf{E}\left[\check{F}(X^{m})\right] = \mathbf{E}\left[\int_{0}^{\infty} \frac{d}{dt} P_{\tau}\check{F}(X^{m}) \,\mathrm{d}\tau\right]$$
$$= \mathbf{E}\left[P_{\tau_{0}}\check{F}(X^{m}) - \check{F}(X^{m})\right] + \mathbf{E}\left[\int_{\tau_{0}}^{\infty} LP_{\tau}\check{F}(X^{m}) \,\mathrm{d}\tau\right],$$

for any $\tau_0 > 0$.

Theorem 4.4. If $p \ge 3$ and X_a belongs to L^p , for any $\tau_0 > 0$, there exists c > 0 such that

(8)
$$\mathbf{E}\left[\int_{\tau_0}^{\infty} LP_{\tau}\check{F}(X^m) d\tau\right] \le c \|X_a\|_{L^{p/2}} m^{-1/2+3\eta} \int_{\tau_0}^{\infty} \frac{e^{-5\tau/2}}{1-e^{-\tau/2}} d\tau.$$

Proof. Let $X_{\neg a}^m = X^m - X_a g_a^m$. Since the X_a 's are independent, $\mathbf{E}\left[\langle \nabla \mathbf{P}, \check{\mathbf{D}}, \check{\mathbf{V}}^m \rangle - X_a^m \rangle\right]$

$$\begin{split} \mathbf{E} \left[\left\langle \nabla P_{\tau} \dot{F}(X^{m}), X^{m} \right\rangle \right] \\ &= \alpha_{m} \mathbf{E} \left[\sum_{a \in \mathcal{A}} X_{a} \left\langle \nabla P_{\tau} \check{F}(X^{m}), g_{a}^{m} \right\rangle \right] \\ &= \alpha_{m} \mathbf{E} \left[\sum_{a \in \mathcal{A}} X_{a} \left\langle \nabla P_{\tau} \check{F}(X^{m}) - \nabla P_{\tau} \check{F}(X_{\neg a}^{m}), g_{a}^{m} \right\rangle \right] \\ &= \alpha_{m}^{2} \mathbf{E} \left[\sum_{a \in \mathcal{A}} X_{a}^{2} \left\langle \nabla^{(2)} P_{\tau} \check{F}(X_{\neg a}^{m}), g_{a}^{m} \otimes g_{a}^{m} \right\rangle \right] \\ &+ \alpha_{m}^{3} \mathbf{E} \left[\sum_{a \in \mathcal{A}} X_{a}^{3} \int_{0}^{1} (1 - r) \left\langle \nabla^{(3)} P_{\tau} \check{F}(X_{\neg a}^{m} + r X_{a} g_{a}^{m}), g_{a}^{m \otimes 3} \right\rangle \, \mathrm{d}r \right], \end{split}$$

according to the Taylor formula. Since $\mathbf{E} \left[X_a^2 \right] = 1$, we have

$$\begin{split} \mathbf{E}\left[LP_{\tau}F(X^{m})\right] \\ &= -\alpha_{m}^{2} \mathbf{E}\left[\sum_{a\in\mathcal{A}}\left\langle \nabla^{(2)}P_{\tau}\check{F}(X_{\neg a}^{m}) - \nabla^{(2)}P_{\tau}\check{F}(X^{m}), \, g_{a}^{m}\otimes g_{a}^{m}\right\rangle\right] \\ &+ \alpha_{m}^{3} \mathbf{E}\left[\sum_{a\in\mathcal{A}}X_{a}^{3}\int_{0}^{1}(1-r) \left\langle \nabla^{(3)}P_{\tau}\check{F}(X_{\neg a}^{m} + r\alpha_{m}X_{a}g_{a}^{m}), \, g_{a}^{m\otimes3}\right\rangle \, \mathrm{d}r\right] \\ &:= -\alpha_{m}^{2} \frac{e^{-3\tau/2}}{\beta_{\tau/2}^{2}} A_{1} + \alpha_{m}^{3} \frac{e^{-5\tau/2}}{\beta_{\tau/2}^{2}} A_{2}. \end{split}$$

In view of (6),

$$A_{1} = \sum_{a \in \mathcal{A}} \mathbf{E} \left[\iint \left[\check{F} \left(X^{m}(\tau, y, z) \right) - \check{F} \left(X^{m}_{\neg a}(\tau, y, z) \right) \right] \right. \\ \left. \times (y, g^{m}_{a})_{H}(z, g^{m}_{a})_{H} \, \mathrm{d}\mathbf{P}^{m}(y) \, \mathrm{d}\mathbf{P}^{m}(z) \right].$$

Since F is Lipschitz continuous, we get

$$|A_1| \le \alpha_m \sum_{a \in \mathcal{A}} B_1^a + \alpha_m \sum_{a \in \mathcal{A}} B_2^a,$$

where

$$\begin{split} B_1^a &= \iint \mathbf{E} \left[|X^m(\tau, y, z) - X^m_{\neg a}(\tau, y, z)|_{W_{\eta, p}} \right] \\ &\times (y, g_a^m)_H(z, g_a^m)_H \, \mathrm{d}\mathbf{P}^m(y) \, \mathrm{d}\mathbf{P}^m(z) \\ B_2^a &= \iint \mathbf{E} \left[\left(\iint_{[0,1]^2} \left| \pi_2 \left(S_2(X^m(\tau, y, z)) - S_2(X^m_{\neg a}(\tau, y, z)) \right)_{s,t} \right|^{p/2} \right] \\ &\mathrm{d}\mu_{\eta, p}(t, s) \right]^{1/p} \right] \times (y, g_a^m)_H(z, g_a^m)_H \, \mathrm{d}\mathbf{P}^m(y) \, \mathrm{d}\mathbf{P}^m(z). \end{split}$$

Since g_a^m has unit norm in $W_{\eta,p}$, we readily have

$$B_1^a \le e^{-\tau} \mathbf{E} [|X_a|] \left(\int |(y, g_a^m)_H| \, \mathrm{d} \mathbf{P}^m(y) \right)^2 \le c \ e^{-\tau} \mathbf{E} [|X_a|^p]^{1/p}.$$

Now then,

$$\begin{split} \left(\pi_2 \check{\mathbf{S}}_2(X^m(\tau, y, z)) - \pi_2 \check{\mathbf{S}}_2(X^m_{\neg a}(\tau, y, z))\right)_{s,t} \\ &= X_a(\tau, y, z) \, \sum_{b \prec a} X_b(\tau, y, z) \, \check{g}^m_{a,b}(s, t). \end{split}$$

Apply Hölder inequality,

$$|B_{2}^{a}| \leq \iint \left(\iint_{[0,1]^{2}} \mathbf{E} \left[\left| X_{a}(\tau, y, z) \sum_{b \prec a} X_{b}(\tau, y, z) \check{g}_{a,b}^{m}(s, t) \right|^{p/2} \right] d\mu_{\eta,p}(t, s) \right)^{1/p} \\ \times |(y, g_{a}^{m})_{H}| |(z, g_{a}^{m})_{H}| d\mathbf{P}^{m}(y) d\mathbf{P}^{m}(z).$$

For y and z fixed, the family $(X_a(\tau, y, z), a \in \mathcal{A})$ satisfies the hypothesis of Lemma 4.3, hence

$$|B_2^a| \le c \iint \mathbf{E} \left[|X_a(\tau, y, z)|^{p/2} \right]^{2/p} |(y, g_a^m)_H| |(z, g_a^m)_H| \, \mathrm{d}\mathbf{P}^m(y) \, \mathrm{d}\mathbf{P}^m(z) \\ \le c \, \|X_a\|_{L^{p/2}}.$$

It follows that for any $a \in \mathcal{A}$, B_1^a and B_2^a are bounded with respect to m by a constant times the p-th moment of X_a to the square. Since $|\mathcal{A}| = dm$, A_1 is bounded by m times this upper-bound. In view of (7), the same kind of computations can be done for A_2 . Combining these upper-bounds, we get the existence of a constant c such that for any $\tau \geq \tau_0$,

$$\left| \mathbf{E} \left[L P_{\tau} \check{F}(X^m) \right] \right| \le c \left\| X_a \right\|_{L^{p/2}} \frac{e^{-5\tau/2}}{\beta_{\tau/2}^2} m \alpha_m^3.$$

Since $\alpha_m = m^{-1/2+\eta}$, the result follows.

Theorem 4.5. For any $\tau_0 > 0$,

$$\left| \mathbf{E} \left[P_{\tau_0} \check{F}(X^m) - \check{F}(X^m) \right] \right| \le c \, \|X_a\|_{L^p} \, m^{-1/2+\eta} \sqrt{1 - e^{-\tau_0}}.$$

Proof.

$$\begin{aligned} \left| \mathbf{E} \left[P_{\tau_0} \check{F}(X^m) - \check{F}(X^m) \right] \right| \\ &\leq \mathbf{E} \left[\int \left| \check{F}(e^{-\tau_0} X^m + \beta_{\tau_0} y) - \check{F}(X^m) \right| \, \mathrm{d}\mathbf{P}^m(y) \right] \\ &\leq \alpha_m \, \mathbf{E} \left[\left| (1 - e^{-\tau_0}) X^m + \beta_{\tau_0} y \right|_{W_{\eta,p}} \, \mathrm{d}\mathbf{P}^m(y) \right] \\ &\quad + \alpha_m \, \mathbf{E} \left[\int \left(\iint_{[0,1]^2} U(X^m)_{s,t} \, \mathrm{d}\mu_{\eta,p}(s,t) \right)^{1/p} \, \mathrm{d}\mathbf{P}^m(y) \right] \end{aligned}$$

where

$$U(X^{m})_{s,t} = \left| \pi_{2} (\check{S}_{2} (\beta_{\tau_{0}}^{2} X^{m} + \beta_{\tau_{0}} y)_{s,t} - \check{S}_{2} (X^{m})_{s,t}) \right|^{p}$$

Thus,

$$\begin{aligned} \left| \mathbf{E} \left[P_{\tau_0} \check{F}(X^m) - \check{F}(X^m) \right] \right| \\ &\leq c \,\beta_{\tau_0} \alpha_m \left(\|X^m\|_{W_{\eta,p}} + \int |y| \, \mathrm{d}\mathbf{P}^m(y) \right) \right) \\ &\leq c \, \|X_a\|_{L^p} \, m^{-1/2+\eta} \sqrt{1 - e^{-\tau_0}}, \end{aligned}$$

according to Corollary 2.3.

References

- R. A. Adams and J. J. F. Fournier, *Sobolev spaces*, second ed., Pure and Applied Mathematics (Amsterdam), vol. 140, Elsevier/Academic Press, Amsterdam, 2003.
- [2] A. D. Barbour, Stein's method for diffusion approximations, Probability Theory and Related Fields 84 (1990), no. 3, 297–322.
- [3] A. D. Barbour and L. H. Y. Chen, An introduction to Stein's method, Lecture Notes Series, vol. 4, National University of Singapore, 2005.
- [4] H. Brézis, Functional analysis, Sobolev spaces and partial differential equations, Springer (New-York), 2011.
- [5] L. Coutin and L. Decreusefond, Stein's method for Brownian approximations, Communications on Stochastic Analysis 7 (2013), no. 3, 349–372.
- [6] L. Decreusefond, Stochastic calculus with respect to Volterra processes, Annales de l'Institut Henri Poincaré (B) Probability and Statistics 41 (2005), 123–149.
- [7] L. Decreusefond, The Stein-Dirichlet-Malliavin method, ESAIM: Proceedings (2015), 11.
- [8] M. D. Donsker, An invariance principle for certain probability limit theorems, Mem. Amer. Math. Soc. 6 (1951).
- [9] R. M. Dudley, *Real analysis and probability*, vol. 74, Cambridge Studies in Advanced Mathematics, no. 3, Cambridge University Press, Cambridge, 2002.
- [10] D. Feyel and A. de La Pradelle, On fractional Brownian processes, Potential Anal. 10 (1999), no. 3, 273–288.
- [11] P. Friz and N. Victoir, Multidimensional stochastic processes as rough paths, Cambridge Studies in Advanced Mathematics, vol. 120, Cambridge University Press, Cambridge, 2010.
- [12] P. Friz and N. Victoir, A variation embedding theorem and applications, Journal of Functional Analysis 239 (2006), no. 2, 631–637.
- [13] J. Lamperti, On convergence of stochastic processes, Transactions of the American Mathematical Society 104 (1962), 430–435.
- [14] I. Nourdin and G. Peccati, Normal Approximations with Malliavin Calculus: From Stein's Method to Universality, vol. 11, Cambridge University Press, 2012.
- [15] S.G. Samko, A.A. Kilbas, and O.I. Marichev, Fractional Integrals and Derivatives, Gordon and Breach Science, jun 1993.

- [16] H.-H. Shih, On Stein's method for infinite-dimensional Gaussian approximation in abstract Wiener spaces, Journal of Functional Analysis 261 (2011), no. 5, 1236–1283.
- [17] D. Williams, Probability with martingales, Cambridge Mathematical Textbooks, vol. 14, Cambridge University Press, Cambridge, 1991.

INSTITUT MATHÉMATIQUE DE TOULOUSE, UNIVERSITÉ P. SABATIER, TOULOUSE, FRANCE *E-mail address*: coutin@math.univ-toulouse.fr

LTCI, TELECOM PARISTECH, UNIVERSITÉ PARIS-SACLAY, 75013, PARIS, FRANCE *E-mail address*: laurent.decreusefond@telecom-paristech.fr