



**HAL**  
open science

## Quantifying environmental impacts associated to sodium alginate extraction from seaweed

Pedro Villanueva-Rey, Paula Perez-Lopez, Stephen K Herbert, Gumersindo D Feijoo, Maria Teresa Moreira

► **To cite this version:**

Pedro Villanueva-Rey, Paula Perez-Lopez, Stephen K Herbert, Gumersindo D Feijoo, Maria Teresa Moreira. Quantifying environmental impacts associated to sodium alginate extraction from seaweed. LCM 2017 (8th international conference on Life Cycle Management), Sep 2017, Luxembourg, Luxembourg. hal-01682195

**HAL Id: hal-01682195**

<https://minesparis-psl.hal.science/hal-01682195v1>

Submitted on 12 Jan 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# **Quantifying environmental impacts associated to sodium alginate extraction from seaweed**

Pedro Villanueva-Rey<sup>1\*</sup>, Paula Pérez-López<sup>1,2</sup>, Stephen K Herbert<sup>3</sup>, Gumersindo Feijoo<sup>1</sup>,  
Maria Teresa Moreira<sup>1</sup>

<sup>1</sup> Department of Chemical Engineering, Institute of Technology, University of Santiago de Compostela. 15782 - Santiago de Compostela, Spain

<sup>2</sup>MINES ParisTech, PSL Research University, Centre Observation, Impacts, Energie (O.I.E.), France

<sup>3</sup>Office of Basic Energy Sciences SC-22.1/Germantown Building, U.S. Department of Energy. 1000 Independence Avenue, SW, Washington, D.C

\* Presenter: Tel.: +34 881816739; E-mail address: [pedro.villanuev@hotmail.com](mailto:pedro.villanuev@hotmail.com)

Seaweeds have been harvested along coast line for different purposes. Potential uses of harvested seaweeds depend on the type, being used either as food for human consumption, feed for aquaculture sector, or fertilizers due to their nutrients content. Kelps —which grow along the West coast of the United States— present a high ratio of sodium alginate within their cell walls, a compound widely used by pharmaceutical and food industries because of its features: viscosity, gel and film formation capacity (FMC Corporation, 2015). This fact explains the increasing interest of extracting alginate from this seaweed at industrial level. During the last five years, several projects aiming at establishing an industry around the

algae *Nereocystis luetkeana* have been developed in the State of Washington, as a possible boost in an economically depressed coast area. In a first phase of this development, a pilot plant is to be created as a spin-off of the University of Wyoming (Laramie). This study focuses on the environmental profile of sodium alginate extraction from *N. luetkeana* in the projected pilot plant by means of life cycle assessment (LCA) methodology. The extraction method assessed was based on a sequence of stages including cutting, soaking with formaldehyde, washing with several solvents, pre-extraction with sulfuric acid and alkaline extraction. The application of Life Cycle Assessment methodology to this process allowed identifying the main hot spots and thereby the inventory items responsible for the environmental impact. The results obtained together with the improvement actions evaluated should be considered for the sake of the optimal valorization of natural resources.

#### References:

Curtiss, P.S., Kreider, J.F., 2009. Algaculture as a Feedstock Source for Biodiesel Fuel: A Life Cycle Analysis 171–179.

EIA. US Energy Information Administration, 2015. Statistics & Analysis: Wyoming -- State Profile and Energy Estimates [WWW Document]. URL <http://www.eia.gov/state> (accessed 23.4.15).

FMC Corporation, 2015. FMC Biopolymer. Alginate [WWW Document]. FMC. URL <http://www.fmcbiopolymer.com/othermarkets/OtherMarkets/Products/Alginate.aspx> (accessed (15.3.14)).

González-López, N., Moure, A., Domínguez, H., 2012. Hydrothermal fractionation of *Sargassum muticum* biomass. *J. Appl. Phycol.* 24, 1569–1578.

Hernández-Carmona, G., McHugh, D.J., López-Gutiérrez, F., 1999. Pilot plant scale extraction of alginates from *Macrocystis pyrifera*. 2. Studies on extraction conditions and methods of separating the alkaline-insoluble residue. *J. Appl. Phycol.* 11, 493–502.

Pérez-López, P., Balboa, E.M., González-García, S., Domínguez, H., Feijoo, G., Moreira, M.T., 2014. Comparative environmental assessment of valorization strategies of the invasive macroalgae *Sargassum muticum*. *Bioresour. Technol.* 161, 137–148.